

# SQUARE-FREE $S$ -MODULES WITH SUPPORT ON A SIMPLICIAL GRAPH AND BRILL–NOETHER THEORY

HENNING LOHNE

**ABSTRACT.** We study square-free  $S$ -modules with support on a simplicial graph, and investigate an analogy between Cohen–Macaulay modules, locally of rank 1, supported on a connected graph and line bundles on a curve. We use the combinatorial structure of the graph to prove a corresponding Riemann–Roch theorem, we study the Jacobian for a connected graph, and we study Brill–Noether theory for 2-connected graphs.

## 1. INTRODUCTION

Let  $S = k[x_1, \dots, x_n]$  be the polynomial ring. K. Yanagawa ([10] and [11]) introduced the notion of square-free  $\mathbf{Z}^n$ -graded  $S$ -modules. Square-free modules is a generalization of square-free monomial ideals. These modules are still of combinatorial nature, as they have support on a simplicial complex. In this paper we study square-free modules whose support is a simplicial graph. Such modules can be thought of as combinatorial analogues to coherent sheaves, or vector bundles if they are Cohen–Macaulay. We study the square-free modules analogue to line bundles on simplicial graphs and compare this with the results from the theory of line bundles on curves. For modules analogue to line bundles we define the degree and global sections, and we show an analogue to the Riemann–Roch theorem (Theorem 3.1) for simplicial graphs. There is another combinatorial Riemann–Roch theorem for graphs, shown by M. Baker and S. Norine in [2]. Our setting is however different, and our result should be thought of as an algebraic combinatorial analogue. Furthermore, we study square-free modules of multi-degree  $(0, \dots, 0)$  with support on a simplicial graph. This family is an analogue to the Jacobian variety of a curve. We show that the dimension of the family of isomorphism classes of such modules equals the genus of the simplicial graph, similar as for the moduli space of line bundles on a curve. We also want to study Brill–Noether theory for modules on a simplicial graph. It turns out that we have to limit this to 2-connected simplicial graphs. We define effective and special square-free modules on such graphs, and we define an analogue to the gonality and Clifford index. We show that the gonality has the same upper bound as in the classical case, and we prove Clifford’s theorem for simplicial graphs. In Remark 5.9, we briefly discuss the possible

---

*Date:* March 9, 2010.

*2000 Mathematics Subject Classification.* Primary: 13F55, 13C14; Secondary: 14H51.

*Key words and phrases.* Cohen–Macaulay modules, Stanley–Reisener rings, Brill–Noether theory.

connection between these numerical invariants and the resolution of graph curves. A similar study has been done by D. Bayer and D. Eisenbud in [3].

The paper is organized as follows: Section 2 contains the preliminaries about square-free  $S$ -modules. We also give a combinatorial description of the canonical module  $k[\Delta]$ .

In Section 3, we describe the analogy between square-free modules and vector bundles. The main theorem of this section is the analogue of the Riemann–Roch theorem. We also give a bound of the degree of an indecomposable square-free module and the existence of some modules of a given degree and a given number of global sections.

In Section 4, we study square-free modules of multi-degree  $(0, \dots, 0)$  with support on a simplicial graph. We show that the dimension of the family of isomorphism classes of such modules are the genus of the simplicial graph, similar as for the moduli space of line bundles on a curve.

In Section 5, we study 2-connected simplicial graphs. We define effective and special square-free modules, and we define an analogue of the gonality and Clifford index. The main results are that the gonality has the same upper bound as in the classical case, and Clifford’s theorem for simplicial graphs.

**Acknowledgments.** I would like to thank Professor Gunnar Fløystad for interesting discussions and valuable suggestions. I also thank Professor Frank-Olaf Schreyer for helpful comments during my stay at the University of Saarland.

## 2. PRELIMINARIES

The most natural square-free  $S$ -modules are square-free monomial ideals and the corresponding quotient rings. These objects are given by the combinatorial structure of simplicial complexes and Stanley–Reisner rings. We recall the following definitions.

Let  $S := k[x_1, x_2, \dots, x_n]$  be the polynomial ring over some field  $k$  and let  $[n] = \{1, 2, \dots, n\}$ . A subset  $F \subseteq [n]$  is called a face. A simplicial complex is a collection of faces  $\Delta$ , such that if  $F \in \Delta$  and  $G \subseteq F$ , then  $G \in \Delta$ . The Stanley–Reisner ideal of the simplicial complex  $\Delta$  is the square-free monomial ideal  $I_\Delta = \langle x^\sigma \mid \sigma \notin \Delta \rangle$  generated by monomials corresponding to non-faces  $\sigma$  of  $\Delta$ . The Stanley–Reisner ring of  $\Delta$  is the quotient ring  $k[\Delta] := S/I_\Delta$ .

**Definition 2.1** (cf. [11] Definition 2.1). *A finitely generated  $\mathbf{Z}^n$ -graded  $S$ -module  $M = \bigoplus_{\mathbf{a} \in \mathbf{Z}^n} M_{\mathbf{a}}$  is said to be square-free, if the following holds*

- i)  *$M$  is  $\mathbf{N}^n$ -graded, that is  $M_{\mathbf{a}} = 0$  for  $\mathbf{a} \notin \mathbf{N}^n$ .*
- ii) *The multiplication map  $M_{\mathbf{a}} \xrightarrow{x_i} M_{\mathbf{a}+\mathbf{e}_i}$  is bijective for all  $\mathbf{a} \in \mathbf{N}^n$  and all  $i \in \text{supp}(\mathbf{a})$ .*

**Theorem 2.2** (cf. [10] Theorem 2.6). *If  $M$  is a square-free  $S$ -module, then so is  $\text{Ext}_S^i(M, \omega_S)$  for all  $i$ .*

Let  $M$  be a  $\mathbf{Z}^n$ -graded  $S$ -module. The Matlis dual of  $M$  is the  $\mathbf{Z}^n$ -graded  $S$ -module  $M^\vee = \text{Hom}_k(M, k)$ . This means that

$$(M^\vee)_{-\mathbf{a}} = \text{Hom}_k(M_{\mathbf{a}}, k),$$

and the multiplication map  $(M^\vee)_{-\mathbf{a}} \xrightarrow{\cdot \mathbf{x}^{\mathbf{b}}} (M^\vee)_{-\mathbf{a}+\mathbf{b}}$  is the transpose of the multiplication map  $M_{\mathbf{a}-\mathbf{b}} \xrightarrow{\cdot \mathbf{x}^{\mathbf{b}}} M_{\mathbf{a}}$ . See [7] and [8, Def. 11.15].

Let  $H_m^i(M)$  denote the local cohomology modules of  $M$ . Then the following holds.

**Theorem 2.3** (Local duality). *For all finitely generated  $\mathbf{Z}^n$ -graded  $S$ -modules  $M$  and all integers  $i$  there exist natural homogeneous isomorphisms*

$$(H_m^i(M))^\vee \cong \text{Ext}_S^{n-i}(M, \omega_S).$$

*Proof.* See [4, Thm. 3.6.19] or [7, Cor. 6.1].  $\square$

The local cohomology modules can be calculated by the cocomplex

$$K^\bullet : 0 \rightarrow K^0 \rightarrow K^1 \rightarrow \cdots \rightarrow K^n \rightarrow 0,$$

where

$$K^i = \bigoplus_{\substack{\sigma \subseteq [n] \\ |\sigma|=i}} M_{\mathbf{x}^\sigma}$$

is the direct sum of the module localized in the square-free monomials of degree  $i$ , and the differential is given on each component as

$$\text{sign}(j, \sigma \cup j) \cdot \text{nat} : M_{\mathbf{x}^\sigma} \rightarrow M_{\mathbf{x}^{\sigma \cup j}}$$

if  $j \notin \sigma$  and 0 otherwise, where  $\text{sign}(j, \sigma) = (-1)^{\#\{i \in \sigma \mid i < j\}}$ . If  $M$  is square-free, then the cohomology  $H^i(K^\bullet) \cong H_m^i(M)$  is easy to calculate. As in [10] we have that if  $M$  is square-free, then

$$(M_{\mathbf{x}^\sigma})_{-\tau} = \begin{cases} M_\sigma & \text{if } \tau \subseteq \sigma \\ 0 & \text{otherwise,} \end{cases}$$

and the natural map  $(M_{\mathbf{x}^\sigma})_{-\tau} \rightarrow (M_{\mathbf{x}^{\sigma \cup \{i\}}})_{-\tau}$  corresponds to the map  $M_\sigma \xrightarrow{\cdot x_i} M_{\sigma \cup \{i\}}$ .

**Definition 2.4.** *Let  $M$  be a square-free Cohen–Macaulay  $S$ -module of dimension  $d$ . The canonical module of  $M$  is defined as the square-free  $S$ -module*

$$\omega_M = \text{Ext}_S^{n-d}(M, \omega_S).$$

**Proposition 2.5.** *If  $M$  is a square-free Cohen–Macaulay module of dimension  $d$  and if  $\tau \subseteq [n]$ , then*

$$\dim_k(\omega_M)_\tau = \sum_{\tau \subseteq \sigma \subseteq [n]} (-1)^{d-|\sigma|} \dim_k M_\sigma.$$

*Proof.* If  $M$  is Cohen–Macaulay of dimension  $d$ , then the local cohomology groups  $H_m^i(M) = 0$  for  $i \neq d$ . So the cocomplex  $K^\bullet$  restricted to degree  $-\tau$  has only cohomology  $(H_m^d(M))_{-\tau}$ , which is isomorphic to  $\text{Hom}_k((\omega_M)_\tau, k)$  by local duality. The result now follows since

$$(K^i)_{-\tau} = \bigoplus_{\substack{|\sigma|=i \\ \tau \subseteq \sigma}} M_\sigma.$$

$\square$

**2.1. The canonical module of  $k[\Delta]$ .** Assume that  $\Delta$  is a Cohen–Macaulay  $(d-1)$ -dimensional simplicial complex, i.e., the Stanley–Reisner ring  $k[\Delta]$  is Cohen–Macaulay of dimension  $d$ . The Matlis dual of the cocomplex  $K^\bullet$  above restricted to positive degrees gives a nice  $\mathbf{Z}^n$ -graded description of  $\omega_{k[\Delta]}$ . The details are written out in [4, Theorem 5.7.3], here we only give the result.

For each  $i = 0, \dots, d$ , let  $G_i$  be the direct sum of the  $k[\Delta]$ -modules  $k[X_1, \dots, X_n]/I_F$  where  $F \in \Delta$ ,  $|F| = i$  and  $I_F = (X_j \mid j \notin F)$ . Let  $\varphi_i : G_i \rightarrow G_{i-1}$  be the map which is

$$(-1)^{j-1} \text{nat} : k[X_1, \dots, X_n]/I_F \rightarrow k[X_1, \dots, X_n]/I_{F'}$$

if  $F = \{v_{i_1}, \dots, v_{i_r}\}$  and  $F' = \{v_{i_1}, \dots, \widehat{v_{i_j}}, \dots, v_{i_r}\}$ , and zero otherwise. Then the complex

$$G_\bullet : 0 \rightarrow G_d \rightarrow G_{d-1} \rightarrow \dots \rightarrow G_1 \rightarrow G_0 \rightarrow 0$$

is the Matlis dual of  $K^\bullet$  restricted to positive degrees, and since  $\Delta$  is Cohen–Macaulay we obtain an exact sequence of  $\mathbf{Z}^n$ -graded  $k[\Delta]$ -modules

$$0 \rightarrow \omega_{k[\Delta]} \rightarrow G_d \rightarrow G_{d-1} \rightarrow \dots \rightarrow G_1 \rightarrow G_0 \rightarrow 0.$$

The description of the canonical module in the long exact sequence above gives rise to a nice description of its square-free grades. Let  $\text{st}_\Delta F$  denote the set of faces of  $\Delta$  containing  $F$ , and let  $\Delta - F$  denote  $\Delta \setminus \text{st}_\Delta F$ . We have that

$$(\omega_{k[\Delta]})_0 \cong \widetilde{H}_{d-1}(\Delta; k) \text{ and}$$

$$(\omega_{k[\Delta]})_F \cong \widetilde{H}_{d-1}(\Delta, \Delta - F; k) \text{ for faces } \emptyset \neq F \in \Delta,$$

and the multiplication map

$$\widetilde{H}_{d-1}(\Delta, \Delta - F; k) \xrightarrow{x_i} \widetilde{H}_{d-1}(\Delta, \Delta - (F \cup \{i\}); k)$$

is the natural map.

**Remark 2.6.** For any face  $F$ , the homology groups of the chain complex above, restricted to degree  $F$ , can also be described in the more common way using the link of a face. Recall that the link of  $F$  in  $\Delta$  is defined as  $\text{lk}_F \Delta := \{G \in \Delta \mid F \cap G = \emptyset \text{ and } F \cup G \in \Delta\}$ . The homology groups above can now be described as  $\widetilde{H}_i(\Delta, \Delta - F; k) \cong \widetilde{H}_{i-|F|}(\text{lk}_F \Delta; k)$ . This is because the chain complexes are the same. The two different descriptions both have their advantages. The first gives a natural description of the multiplication map, while the second is a combinatorial description.

We use these homology groups to give a characterisation of 2-CM simplicial complexes. Recall that a Cohen–Macaulay simplicial complex  $\Delta$  is said to be doubly Cohen–Macaulay or 2-CM if  $\Delta - p$  is Cohen–Macaulay of the same dimension as  $\Delta$  for all vertices  $p \in \Delta$ .

**Proposition 2.7.** *Let  $\Delta$  be a simplicial complex of dimension  $d-1$ . Then the following are equivalent.*

1.  $\Delta$  is 2-CM.
2.  $\widetilde{H}_i(\Delta; k) = 0$  for  $0 \leq i \leq d-2$  and  $\widetilde{H}_i(\Delta - F; k) = 0$  for  $0 \leq i \leq d-2$  and any face  $F \in \Delta$ .

*Proof.* Since the homology groups  $\tilde{H}_i(\Delta; k)$  and  $\tilde{H}_i(\Delta, \Delta - F; k)$  are the generators for the module  $\text{Ext}_S^{n-i-1}(k[\Delta], \omega_S)$ , and because of the long exact sequence

$$\cdots \rightarrow \tilde{H}_i(\Delta; k) \rightarrow \tilde{H}_i(\Delta, \Delta - F; k) \rightarrow \tilde{H}_{i-1}(\Delta - F; k) \rightarrow \tilde{H}_{i-1}(\Delta; k) \rightarrow \cdots$$

we see that  $\Delta$  is CM if and only if  $\tilde{H}_i(\Delta; k) = 0$  for  $0 \leq i \leq d-2$  and  $\tilde{H}_i(\Delta - F; k) = 0$  for  $0 \leq i \leq d-3$  and all faces  $\emptyset \neq F \in \Delta$ . So it is enough to show that  $\Delta$  is 2-CM if and only if  $\Delta$  is CM and that  $\tilde{H}_{d-2}(\Delta - F; k) = 0$  for any face  $F \in \Delta$ .

Suppose that  $\Delta$  is 2-CM. We will show that  $\tilde{H}_{d-2}(\Delta - F; k) = 0$  for all faces  $F \in \Delta$  using induction on the dimension of  $F$ . If  $F = p$  is just a vertex, then  $\tilde{H}_{d-2}(\Delta - p; k) = 0$  since  $\Delta - p$  is CM. Now suppose that  $G = F \cup p$  is a face. Let  $A = \Delta - F$  and  $B = \Delta - p$ . Then  $A \cup B = \Delta - G$  and  $A \cap B = \Delta - p - F$ . We therefore have a Mayer-Vietoris sequence

$$\begin{aligned} \cdots \rightarrow \tilde{H}_{d-2}(\Delta - F; k) \oplus \tilde{H}_{d-2}(\Delta - p; k) &\rightarrow \tilde{H}_{d-2}(\Delta - G; k) \rightarrow \\ &\tilde{H}_{d-3}(\Delta - p - F; k) \rightarrow \cdots \end{aligned}$$

The homology groups on the left side are trivial because of the induction hypothesis and the homology group on the right side is trivial because  $\Delta - p$  is CM of dimension  $d-1$  so it follows that  $\tilde{H}_{d-2}(\Delta - G; k) = 0$ .

Suppose that  $\Delta$  is CM and that  $\tilde{H}_{d-2}(\Delta - F; k) = 0$  for all faces  $F \in \Delta$ . We have that  $\tilde{H}_i(\Delta - p; k) = 0$  for  $0 \leq i \leq d-2$ . Let  $p \in \Delta$  be a vertex and  $F \in \Delta$  a face not containing  $p$ . As above, we get a Mayer-Vietoris sequence

$$\cdots \rightarrow \tilde{H}_{i+1}(\Delta - G; k) \rightarrow \tilde{H}_i(\Delta - p - F; k) \rightarrow \tilde{H}_i(\Delta - p; k) \oplus \tilde{H}_i(\Delta - F; k) \rightarrow \cdots,$$

where  $G$  is the face  $F \cup p$  if this is a face in  $\Delta$ , otherwise, we just replace  $\Delta - G$  above with  $\Delta$ . For  $0 \leq i \leq d-3$ , the homology group on the left side is trivial because of the assumption and the fact that  $\Delta$  is CM, and the homology groups on the right side are trivial because  $\Delta$  is CM, so  $\tilde{H}_i(\Delta - p - F; k) = 0$  for  $0 \leq i \leq d-3$  and it follows that  $\Delta - p$  is CM for all vertices  $p \in \Delta$ , hence  $\Delta$  is 2-CM.  $\square$

This characterisation of 2-CM simplicial complexes has the following corollary, which we will use later. The only if part is well known. For more details see [1], [4] and [9]. We include a proof of the equivalence for completeness.

**Corollary 2.8.** *Let  $\Delta$  be a CM simplicial complex of dimension  $d-1$ . Then  $\Delta$  is 2-CM if and only if  $\omega_{k[\Delta]}$  is generated by  $(\omega_{k[\Delta]})_0 \cong \tilde{H}_{d-1}(\Delta; k)$ .*

*Proof.* Consider the long exact sequence

$$0 \rightarrow \tilde{H}_{d-1}(\Delta - F) \rightarrow \tilde{H}_{d-1}(\Delta) \xrightarrow{\cdot \mathbf{x}^F} \tilde{H}_{d-1}(\Delta, \Delta - F) \rightarrow \tilde{H}_{d-2}(\Delta - F) \rightarrow 0.$$

$\omega_{k[\Delta]}$  is generated by  $\tilde{H}_{d-1}(\Delta; k)$  if and only if  $\cdot \mathbf{x}^F$  is surjective for every  $F$ . That is, if and only if  $\tilde{H}_{d-2}(\Delta - F) = 0$  for every face  $F$ .  $\square$

## 3. THE RIEMANN–ROCH THEOREM

The annihilator of a square-free  $S$ -module  $M$  is a square-free monomial ideal. Since square-free monomial ideals corresponds to simplicial complexes,  $M$  can be considered as a module over the Stanley–Reisner ring  $k[\Delta] = k[x_1, \dots, x_n]/\text{ann}(M)$ . The study of square-free Cohen–Macaulay modules are therefore the same as the study of maximal Cohen–Macaulay modules over Stanley–Reisner rings  $k[\Delta]$ , with support on all of  $k[\Delta]$ . In the rest of this paper we will study square-free Cohen–Macaulay modules with support on a simplicial graph. These modules are interesting because there is an analogy between line bundles on an algebraic curve and square-free CM  $S$ -modules with support on a simplicial graph.

A square-free  $S$ -module can be described as follows. For each face  $F \subseteq [n]$  we have a  $k$ -vector space  $M_F$ , and for each face  $F \subseteq G$  we have a  $k$ -linear map  $\varphi_{FG} : M_F \rightarrow M_G$  satisfying  $\varphi_{FF} = \text{id}_{M_F}$  and if  $F \subseteq G \subseteq H$ , then  $\varphi_{GH} \circ \varphi_{FG} = \varphi_{FH}$ .

When  $M$  is CM, or more generally when  $\text{depth}(M) \geq 2$ , this description of a module  $M$  on  $k[\Delta]$  is a combinatorial analogy of the sheaf  $\widetilde{M}$  on  $\text{Proj } k[\Delta]$ . This is because of the natural graded isomorphism  $M \cong \bigoplus_{\tau} H^0(\text{Proj } k[\Delta], \widetilde{M}(\tau))$  [6, Appendix A4]. Furthermore, when  $M_F \cong k$  for all edges  $F \in \Delta$ , it is an analogy of a line bundle. In this case we say that  $M$  is locally of rank 1. Inspired by this, we define  $l(M) := \dim_k M_{\emptyset}$  as an analogy for the global sections. Furthermore, we define the multi-degree of  $M$  as the vector  $\mathbf{d} \in \mathbf{Z}^n$ , such that  $d_i = \dim_k M_{\{i\}} - 1$ , and the degree of  $M$  as  $\deg(M) := \sum_i d_i$ . This definition of the degree of a module is an analogy to the degree of a line bundle.

Let  $\Delta$  be a connected simplicial complex of dimension 1 on the vertex set  $[n]$ , and set  $v := \#\{\text{vertices in } \Delta\}$  and  $e := \#\{\text{edges in } \Delta\}$ . We define the genus of  $\Delta$  as

$$g(\Delta) := l(\omega_{k[\Delta]}) = \dim_k (\omega_{k[\Delta]})_0 = \dim_k \widetilde{H}_1(\Delta; k) = 1 - v + e.$$

**Theorem 3.1** (Riemann–Roch). *Let  $\Delta$  be a connected simplicial graph, and let  $M$  be a square-free Cohen–Macaulay module with support on  $\Delta$ , locally of rank 1. Then the following formula holds.*

$$l(M) - l(\omega_M) = 1 + \deg(M) - g.$$

*Proof.* Using Proposition 2.5, we can calculate the left hand side as

$$\sum_{|\sigma|=1} M_{\sigma} - \sum_{|\tau|=2} M_{\tau} = (\deg(M) + v) - e = \deg(M) + 1 - g.$$

□

Using the Riemann–Roch formula, it is easy to see that  $\deg(\omega_M) = 2g - 2 - \deg(M)$ . Since the degree is obviously bounded below by  $\deg(M) \geq -v$  and  $\omega_M$  is a square-free Cohen–Macaulay module whenever  $M$  is, we get that the degree is also bounded above by  $\deg(M) \leq 2g - 2 + v$ . For a given simplicial graph, it is possible to give a better bound for the degree. It is not so difficult to see that for a module to be indecomposable, it must have non-zero vector spaces in the vertices in a vertex cover  $C$  of  $\Delta$ , where  $\Delta|_C$  is connected. This leads to the following proposition.

**Proposition 3.2.** *Let  $\Delta$  be a connected simplicial graph, and let  $M$  be an indecomposable square-free  $S$ -module with support on  $k[\Delta]$ . Then*

$$-s \leq \deg(M) \leq 2g - 2 + s,$$

where

$$s = \max \{ |\sigma| \mid \sigma \subseteq [n], \dim(\Delta|_\sigma) = 0 \text{ and } \Delta|_{\sigma^c} \text{ is connected} \}.$$

*Proof.* Since the upper bound is the dual of the lower bound, it is enough to show that the lower bound holds. Suppose that  $\deg(M) = -k$ . Then there are at least  $k$  vertices in  $\Delta$  where  $M_v = 0$ . Let  $\tau \subseteq [n]$  be the subset that corresponds to these vertices. Since  $M$  is indecomposable,  $\Delta|_{\tau^c}$  must be connected and there can not be any edges in  $\Delta|_\tau$ , so  $\dim(\Delta|_\tau) = 0$ . It therefore follows from the definition of  $s$  in the proposition that  $-s \leq -k$ .

Let  $\sigma \subseteq [n]$  be a subset such that  $|\sigma| = s$ ,  $\dim(\Delta|_\sigma) = 0$  and  $\Delta|_{\sigma^c}$  is connected. Then we can construct an indecomposable module  $M'$  with  $\deg(M') = -s$ . Let  $M'$  be the module where  $M'_v = 0$  for all vertices  $v \in \sigma$ ,  $M'_v \cong k$  for all vertices  $v \in \sigma^c$  and  $M'_e \cong k$  for all edges, with multiplication map  $\varphi_{ve} = 1$  for all vertices  $v \in \sigma^c$ . Then since  $\dim(\Delta|_\sigma) = 0$  all edges are in  $\Delta|_{\sigma^c}$ , which is connected, so  $M'$  is indecomposable. It also follows from the construction that  $\deg(M') = -s$ .  $\square$

We also have the following result, which is an analogy to a well-known fact from algebraic geometry.

**Proposition 3.3.** *Let  $\Delta$  be a connected simplicial graph of genus  $g$ . Then for any  $0 \leq d \leq g$ , there exists an indecomposable CM module  $M$  with support on  $k[\Delta]$ , locally of rank 1, with  $l(M) = 1$ ,  $\deg(M) = d$  and multi-degree  $\geq (0, 0, \dots, 0)$ .*

*Proof.* If  $\Delta$  is as above, then  $\Delta = \Gamma \cup \{e\}$  for some edge  $e \in \Delta$  and some connected simplicial graph  $\Gamma$  of genus  $g - 1$ . By induction, we may assume that the proposition holds for  $\Gamma$ . Let  $M'$  be such a module on  $k[\Gamma]$  with  $\deg(M') = d' \leq g - 1$ . Form the module  $M''$  on  $\Delta$  such that  $M''|_\Gamma = M'$  and  $M''_e \cong k$ , with  $k$ -linear maps  $\varphi_{ve} = 0$  and  $\varphi_{we}$  defined such that  $\varphi_{we} \circ \varphi_{\emptyset w} = 1_e$ . Let  $M$  be the module such that  $M_\sigma = M''_\sigma$  if  $\sigma \neq v$  and  $M_v = M''_v \oplus k\{1_b\}$ , and let the  $k$ -linear maps of  $M$  be the same as for  $M''$ , except that  $\varphi_{ve}(1_b) = 1_e$ , and  $\varphi_{\emptyset v} = 1_b$ . We get that  $\deg(M) = d' + 1$ , and by the construction of  $M$  we see that the number of “global sections” is  $l(M) = l(M') = 1$ . If we analyze the cocomplex  $K^\bullet$  restricted to the negative square-free degrees, we see that  $M$  is CM if and only if the maps  $\varphi_v (= \bigoplus_{v \in e_i} \varphi_{ve_i})$  and  $\bigoplus \varphi_v$  are injective. This holds since  $M'$  is CM, so  $M$  is also CM.

The case where  $d = 0$  follows easily, since we can just choose  $M = k[\Delta]$ .  $\square$

#### 4. THE JACOBIAN

We study the moduli space of isomorphism classes of square-free CM  $S$ -modules, with support on a simplicial graph, of multi-degree  $(0, 0, \dots, 0)$ . This space is not as nice as the Jacobi variety of an algebraic curve. However, if we restrict to “non-degenerate” modules we can give the space an algebraic structure with dimension equal the genus of the graph.

**Lemma 4.1.** *If  $\Delta$  is a connected simplicial graph of genus 0, and  $M$  is any square-free indecomposable module (possibly not CM) of multi-degree  $(0, 0, \dots, 0)$  with support on  $\Delta$ , locally of rank 1, then  $M$  is isomorphic to a module with  $k$ -linear maps  $\varphi_{ve} = 1$  if  $v \in e$  and 0 otherwise.*

*Proof.* We do induction on the number of vertices on  $\Delta$ . Suppose  $\Delta = \Gamma \cup v \cup e$ , where  $e = vw$  is an edge connecting  $v$  and  $\Gamma$ . By induction it is enough to show that  $\varphi_{we} = 1$  and  $\varphi_{ve} = 1$ . However, since both  $\varphi_{we}$  and  $\varphi_{ve}$  are non-zero (if one of them is zero then  $M$  is decomposable) we can first find a basis of  $M_e$  such that  $\varphi_{we} = 1$ , and next find a basis for  $M_v$  such that  $\varphi_{ve} = 1$ . The same technique can be used to show the case where  $\Delta$  has only one edge. This completes the proof.  $\square$

**Proposition 4.2.** *Let  $\Delta$  be the boundary of an  $n$ -gon. The family of indecomposable CM modules over  $k[\Delta]$ , locally of rank 1, with multi-degree  $(0, 0, \dots, 0)$  is parametrized by a union of  $n$   $\mathbf{P}_k^1$ 's, where we identify the same points from the different  $\mathbf{P}_k^1$ 's if they are not  $(0, 1)$  or  $(1, 0)$ . In other words, its parametrized by a  $\mathbf{P}_k^1$ , with the points  $(0, 1)$  and  $(1, 0)$   $n$ -doubled.*

*Proof.* If  $M$  is indecomposable, then there exists an edge  $e$  in  $\Delta$  such that if we set  $\Gamma := \Delta - \{e\}$ , then  $M|_\Gamma$  is indecomposable. Since  $\Delta$  is the boundary of a polygon,  $\Gamma$  is a connected simplicial graph of genus 0. This means that we may assume that  $M|_\Gamma$  has the  $k$ -linear maps as described in the lemma above.  $M$  is therefore determined by the two  $k$ -linear maps  $\varphi_{ve}$  and  $\varphi_{we}$ , where  $e = vw$ . For any basis of  $M_e$ , these maps are determined by a pair  $(s, t) \in k^2$ , and  $M$  is decomposable if and only if  $s = t = 0$ . Since any basis change of  $M_e$  is multiplication by a non-zero element of  $k$ , it follows that the maps are determined, up to isomorphism, by an element  $(s, t) \in \mathbf{P}_k^1$ .

Suppose that edge chosen is not unique, and that there is another edge  $e'$  such that  $\Gamma' := \Delta - \{e'\}$  is indecomposable. Since  $\Gamma'$  also satisfies the condition for the lemma above, we must also have that  $M|_{\Gamma'}$  is isomorphic to a module with the  $k$ -linear maps  $\varphi_{v_i e_j} = 1$ . This can only happen if both  $s \neq 0$  and  $t \neq 0$ . We may therefore change the basis for the appropriate vector spaces, one by one, to obtain a situation where the pair  $\varphi_{v' e'}$  and  $\varphi_{w' e'}$  is determined by the same pair  $(s, t)$  and all the other  $k$ -linear maps is 1. This explains the identification of points in the parametrization. The edge is unique if either  $(s, t) = (1, 0)$  or  $(0, 1)$ .

By analyzing the complex  $K^\bullet$  in the negative square-free degrees, it follows that  $M$  is CM if all the maps  $\varphi_v := \bigoplus_{v \in e_i} \varphi_{ve_i}$  are injective and the image  $\bigoplus \varphi_{\emptyset v_i}(M_\emptyset)$  equals the kernel of  $\bigoplus \varphi_v$ . It therefore follows that any module created above is CM if  $M_\emptyset$  and  $\varphi_{\emptyset v_i}$  are chosen properly. It can be shown that  $l(M) = 1$  if  $(s, t) = (1, 1)$ , and  $l(M) = 0$  otherwise.  $\square$

If we only consider the modules where all the maps  $\varphi_{ve}$  are non-zero, we get from the proof of the proposition above that the family of the modules above is parametrized by  $\mathbf{A}_k^1 \setminus \{0\}$ .

**Proposition 4.3.** *Let  $\Delta$  be a simplicial graph of genus  $g \geq 1$ . The family of indecomposable CM modules over  $k[\Delta]$ , locally of rank 1, with multi-degree  $(0, 0, \dots, 0)$  and where all the maps  $\varphi_{ve}$  are non-zero is parametrized by  $(\mathbf{A}_k^1 \setminus \{0\})^g$ .*

*Proof.* If  $g = 1$  then  $\Delta$  is the boundary of an  $n$ -gon union a finite number of trees. The result follows from the proposition and lemma above since the two constructions glue together. Suppose now that the claim holds for all graphs of genus  $g \leq k$ , and suppose that  $\Delta$  is a graph of genus  $k + 1$ . Then  $\Delta = \Gamma \cup \{e\}$ , where  $\Gamma$  is a graph of genus  $k$  and  $e = vw$  is an edge. Since any square-free module with support on  $\Delta$  is given as a square-free module on  $\Gamma$  and two maps  $\varphi_{ve}$  and  $\varphi_{we}$ , the result follows by the induction hypothesis and the fact that there is an  $\mathbf{A}_k^1 \setminus \{0\}$  parametrizing the possible choices of  $\varphi_{ve}$  and  $\varphi_{we}$ .  $\square$

## 5. BRILL–NOETHER THEORY

As we have seen, there is an analogy between line bundles on a given curve and square-free Cohen–Macaulay modules with support on a given graph. We investigate this further by defining effective and special modules as an analogue of effective special divisors. The corresponding “Brill–Noether” theory for square-free Cohen–Macaulay modules has some similarities and some differences from the classical theory.

**Definition 5.1.** *Let  $\Delta$  be a Cohen–Macaulay simplicial graph. An indecomposable square-free CM module  $M$  is said to be effective if  $M$  has a submodule isomorphic to  $k[\Delta]$ , and said to be special if  $M \subseteq \omega_{k[\Delta]}$ .*

A simplicial graph is CM if and only if it is connected, and 2-CM if and only if it is 2-connected. From Proposition 2.8, we know that the canonical module  $\omega_{k[\Delta]}$  is generated by the cycles  $\tilde{H}_1(\Delta; k)$  if and only if  $\Delta$  is 2-connected. It is therefore natural to study special modules of 2-connected simplicial graphs.

Some of the following results are using that the field  $k$  is infinite. We are therefore assuming that  $\text{char}(k) = 0$  for the rest of this section.

**Lemma 5.2.** *Let  $\Delta$  be a 2-connected simplicial graph. Then  $\omega_{k[\Delta]}$  has a submodule  $N \cong k[\Delta]$  such that any square-free module  $N \subseteq M \subseteq \omega_{k[\Delta]}$  is indecomposable, and therefore effective.*

*Proof.* Let  $\{s_1, \dots, s_g\}$  be a basis for  $\tilde{H}_1(\Delta; k)$ , and let  $u_N = c_1 s_1 + \dots + c_g s_g$  be a general element in  $\tilde{H}_1(\Delta; k)$ . Then  $\text{Supp}(u_N) = \Delta$ . Let  $N$  be the submodule of  $\omega_{k[\Delta]}$  generated by  $u_N$  and suppose that  $M$  is a module  $N \subseteq M \subseteq \omega_{k[\Delta]}$ . Suppose that  $M = M' \oplus M''$  is decomposable. It is not possible to have  $u_N \in M'$ , since any non-zero element  $m \in M''$  has support on a facet  $F$  and since  $M_F \cong k$ . Suppose that  $u_N = m_1 + m_2$ . We have that  $m_1 = a_1 s_1 + \dots + a_g s_g$  and  $m_2 = b_1 s_1 + \dots + b_g s_g$ , where  $a_i + b_i = c_i$ . For the same reason as above, we must have that  $\dim(\text{Supp}(m_1) \cap \text{Supp}(m_2)) < 1$ . This means that  $a_i = 0$  or  $a_i = c_i$ . So we may assume that  $m_1 = c_1 s_1 + \dots + c_h s_h$  and  $m_2 = c_{h+1} s_{h+1} + \dots + c_g s_g$ . Since  $u_N$  is a general element, we have that  $\Gamma_1 := \text{Supp}(m_1) = \text{Supp}(s_1, \dots, s_h)$  and  $\Gamma_2 := \text{Supp}(m_2) = \text{Supp}(s_{h+1}, \dots, s_g)$ . Since  $\Delta$  is connected, we must have that  $\Gamma_1 \cap \Gamma_2 = \{v_1, \dots, v_l\}$ , and since  $\Delta$  is 2-connected, we must have that  $l \geq 2$ . Furthermore, since  $\Delta$  is 2-connected there exists a cycle  $s \in \tilde{H}_1(\Delta; k)$  such that  $v_1, v_2 \in \text{Supp}(s)$  and  $\text{Supp}(s)$  intersects both  $\Gamma_1$  and  $\Gamma_2$  in facets.

We construct the cycle this way: Let  $w_1$  be a vertex adjacent to  $v_1$  in  $\Gamma_1$  and  $w_2$  be a vertex adjacent to  $v_1$  in  $\Gamma_2$ . Since  $\Delta$  is 2-connected,  $\Delta - \{v_1\}$  is connected, so there are cycle-free paths  $\gamma_1$  from  $w_1$  to  $v_2$  and  $\gamma_2$  from  $w_2$  to  $v_2$ . Let  $s \in \tilde{H}_1(\Delta; k)$  be a cycle corresponding to the cycle  $\gamma_1$  and  $\gamma_2$  connected in  $v_1$ . Now  $s = d_1 s_1 + \dots + d_g s_g$ . But then  $s|_{\Gamma_1} = d_1 s_1 + \dots + d_h s_h$  is a cycle. This is not possible, since the path  $\gamma_1$  was chosen to be without cycles.  $\square$

**Definition 5.3.** Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 2$ . A special and effective module  $M$  satisfying  $l(M) = r + 1$  and  $\deg(M) = d$  is called a  $g_d^r$ . We define the gonality of a graph to be

$$\text{gon}(\Delta) = \min \{k \mid \Delta \text{ possesses a } g_k^1\}.$$

For curves, it is well known that the gonality lies between 2 and  $\left\lfloor \frac{g+3}{2} \right\rfloor$ . As we will see below, the same holds for a 2-connected simplicial graph.

**Lemma 5.4.** Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 2$ . Then the gonality of  $\Delta$  is less than or equal to the girth (i.e. the length of the shortest closed path) of  $\Delta$ .

*Proof.* From Proposition 5.2, we can choose  $N \subseteq \omega_k[\Delta]$  such that  $N \cong k[\Delta]$ . Choose the cycle  $u_N \in N$  and a cycle  $c$  of length  $w = \text{girth}(\Delta)$ . Then the submodule  $M \subseteq \omega_k[\Delta]$  generated by  $u_N$  and  $c$  is a  $g_d^1$ , where  $d \leq w$  since  $\text{Supp}(u_N) \cap \text{Supp}(c)$  contains no more than  $w$  vertices.  $\square$

**Lemma 5.5.** Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 4$  and minimum valency  $k \geq 3$ . Then the girth of  $\Delta$  is  $\leq \left\lfloor \frac{g+3}{2} \right\rfloor$  for all graphs except three special cases.


*Proof.* According to [5], Theorem 11.11.3, we have that the number of vertices of  $\Delta$ , denoted by  $v$ , is bounded by

$$2g - 2 \geq v \geq 1 + k \frac{(k-1)^{\left\lfloor \frac{w-1}{2} \right\rfloor} - 1}{k-2}.$$


Suppose that  $w \geq \left\lfloor \frac{g+5}{2} \right\rfloor$ , then the inequality above has only finitely many solutions. The possible solutions are given as follows.

| $g$ | $v$   | $w$ |
|-----|-------|-----|
| 4   | 4–6   | 4   |
| 6   | 10    | 5   |
| 7   | 10–12 | 6   |
| 8   | 10–14 | 6   |
| 12  | 22    | 8   |

However, the lower bound can be sharpened. If  $w = 4$  then  $\Delta$  must have a subgraph as in the picture below.



Since  $\Delta$  has minimum valency  $k \geq 3$ , it is clear that it has a subgraph consisting of vertices 1–4. But furthermore, since the vertex 4 has valency  $\geq 3$ , and since the girth is 4 there must exist two other vertices adjacent to 4. Likewise, if  $w = 6$  and  $\Delta$  has minimum valency  $k \geq 3$  then  $\Delta$  must contain a subgraph like the following figure.



The reason for this is the following.  $\Delta$  must contain a vertex, so 1 is in  $\Delta$ . Since  $\Delta$  has minimum valency  $k \geq 3$ , there has to be three vertices adjacent to 1, call them 2, 3 and 4. The vertex 2 must also be adjacent to three vertices. But since the girth  $w = 6$  it is not adjacent to the vertex 2 or 3. Therefore it must be adjacent to two new vertices, call them 5 and 6. The same logic applies to the vertices 3, 4, 5 and 6, and it is clear that  $\Delta$  contains a subgraph as in the figure.

For  $w = 8$ ,  $\Delta$  must also contain a subgraph as above. However, it is not so difficult to see, using the same argument as above, that it is impossible to construct a graph with  $w = 8$  and  $k \geq 3$  with only 8 more vertices. Therefore most of the solutions from the inequality above can not correspond to a graph. The only graphs satisfying the inequality in the Lemma that can exist have the following numerical data.

| $g$ | $v$ | $w$ |
|-----|-----|-----|
| 4   | 6   | 4   |
| 6   | 10  | 5   |
| 8   | 14  | 6   |

There is exactly one graph for each of the cases above. The first is  $K_{3,3}$ , the second is the Peterson graph and the last is the Heawood graph.  $\square$

For any 2-connected graph we would like to find a “canonical” model for it. Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 2$ . If  $v \in G$  is a vertex of valency 2 and  $e_1 = vw_1$  and  $e_2 = vw_2$  are the edges that meet  $v$  and  $e' = w_1w_2$  is not an edge in  $\Delta$ . Then we construct a new graph

$\Delta' = (\Delta - v) \cup \{e' = w_1w_2\}$ . If we iterate this process, we end up with a graph we call  $\tilde{\Delta}$ . It is easy to verify that  $\tilde{\Delta}$  has the same topological structure as  $\Delta$ . We will say that a simplicial graph  $\Delta$  is reduced if  $\Delta = \tilde{\Delta}$ .


**Lemma 5.6.** *Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 2$ . Then  $\text{gon}(\Delta) = \text{gon}(\tilde{\Delta})$ , and  $\tilde{\Delta}$  is either hyperelliptic, i.e. 2-gonal, or has minimal valency  $k \geq 3$ .*

*Proof.* Let  $M$  be an effective module on  $\Delta$ , and let  $v$  be a vertex as described above. Then  $k[\Delta]_v \subseteq M_v \subseteq (\omega_k[\Delta])_v$  but  $k[\Delta]_v \cong k \cong (\omega_k[\Delta])_v$  since  $v$  has valency 2. This means that  $M_v \cong k$ , and it does not contribute to the degree of  $M$ . That means that for any effective module on  $\Delta$  we can construct an effective module  $M'$  on  $\Delta'$  such that  $M'$  has the same numerical invariants as  $M$ . Iterating this process we get an effective module  $\tilde{M}$  on  $\tilde{\Delta}$  with the same numerical invariants as  $M$ . It is also easy to see that any effective module  $N$  on  $\tilde{\Delta}$  can be obtained from an effective module coming from  $\Delta$ . This proves the first part.


For the other part, we notice that either  $\tilde{\Delta}$  has minimal valency  $k \geq 3$  or there exists a vertex  $v$  of valency 2 such that if  $e_1 = vw_1$  and  $e_2 = vw_2$  are the edges that meet  $v$ , then  $e' = w_1w_2$  is an edge in  $\tilde{\Delta}$ . That means that there exists a cycle  $s \in \tilde{H}_1(\tilde{\Delta}; k)$  such that  $\text{Supp}(s) = e_1 \cup e_2 \cup e'$ . Since  $\Delta$  is 2-connected we can find a submodule  $N \subseteq \omega_k[\Delta]$ , isomorphic to  $k[\Delta]$ , such that any module in between is effective. Let  $u_N$  be the generator of  $N$ , then  $M = (u_N, s)$  is a  $g_d^1$  where  $d \leq 3$  since  $\text{Supp}(u_N) \cap \text{Supp}(s)$  contains 3 vertices. Furthermore,  $d \leq 2$  since  $v$  has valency 2, and  $d \geq 2$  can either be seen directly, or by using Proposition 5.8 below. Hence  $\tilde{\Delta}$  has gonality 2, that is  $\tilde{\Delta}$  is hyperelliptic.  $\square$

**Corollary 5.7.** *Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 2$ . Then the gonality of  $\Delta$  is  $\leq \left\lfloor \frac{g+3}{2} \right\rfloor$*

*Proof.* It is enough to show that there exists a  $g_{\left\lfloor \frac{g+3}{2} \right\rfloor}^1$  on the three special graphs above. The graph  $K_{3,3}$  can be drawn like this:



We notice the two subgraphs




Let  $s_1 \in \tilde{H}_1(\tilde{\Delta}; k)$  be a cycle with support on  $\Gamma_1$ . We claim that there exists a cycle  $s_2 \in \tilde{H}_1(\tilde{\Delta}; k)$  with support on  $\Gamma_2$  such that the module  $M$  generated by  $s_1$  and  $s_2$  is a  $g_3^1$ . First of all we notice that for any  $s_2$  with support on  $\Gamma_2$  we have that  $M$  is effective since  $\text{Supp}(s_1, s_2) = \Delta$ , and indecomposable


since  $\dim_k \Gamma_1 \cap \Gamma_2 = 1$ . Next, we show that we can choose  $s_2$  such that the restrictions of  $s_1$  and  $s_2$  are equal in the vertex  $v$ . This follows if we show that we can choose  $s_2$  such that the restrictions of  $s_1$  and  $s_2$  are equal in the three edges of  $v$ . Since  $s_1$  and  $s_2$  are cycles it is enough to show that the restrictions are equal in any two of the edges of  $v$ . But it is clear that we can choose such a  $s_2$  since  $\dim_k \tilde{H}_1(\Gamma_2; k) = 2$ . We now have that

$$\deg(M) = \sum_i (\dim_k (\text{Span}\{s_1|_{v_i}, s_2|_{v_i}\}) - 1) = 3.$$

The Peterson graph is given as:




We notice the two subgraphs




Let  $s_1 \in \tilde{H}_1(\Delta; k)$  be a cycle with support on  $\Gamma_1$ . As above, we claim that there exists a cycle  $s_2 \in \tilde{H}_1(\Delta; k)$  with support on  $\Gamma_2$  such that the module  $M$  generated by  $s_1$  and  $s_2$  is a  $g_4^1$ . For any  $s_2$  with support on  $\Gamma_2$  we have that  $M$  is effective and indecomposable for the same reasons as above. Next, we show that we can choose  $s_2$  such that the restrictions of  $s_1$  and  $s_2$  are equal in the vertices  $v$  and  $v'$ . As above, this follows if we can choose  $s_2$  such that the restrictions of  $s_1$  and  $s_2$  are equal on any two of the edges of  $v$  and any two of the edges of  $v'$ . Again, this follows since  $\dim_k \tilde{H}_1(\Gamma_2; k) = 3$ . Similarly as above, we now get that  $\deg(M) = 4$ .

The Heawood graph is given as:



We notice the two subgraphs



Let  $s_1 \in \tilde{H}_1(\Delta; k)$  be a cycle with support on  $\Gamma_1$ . As above, we claim that there exists a cycle  $s_2 \in \tilde{H}_1(\Delta; k)$  with support on  $\Gamma_2$  such that the module  $M$  generated by  $s_1$  and  $s_2$  is a  $g_4^1$ . For any  $s_2$  with support on  $\Gamma_2$  we have that  $M$  is effective and indecomposable for the same reasons as above. Next, we show that we can choose  $s_2$  such that the restrictions of  $s_1$  and  $s_2$  are equal in the vertices  $v, v'$  and  $v''$ . As above, this follows if we can choose  $s_2$  such that the restrictions of  $s_1$  and  $s_2$  are equal on any two of the edges of  $v, v'$  and  $v''$ . Again, this follows since  $\dim_k \tilde{H}_1(\Gamma_2; k) = 4$ . Similarly as above, we now get that  $\deg(M) = 5$ .  $\square$

One may ask which gonalitys that occur in the bound. It is well-known that in the moduli space of curves of genus  $g$ , there exist curves of gonality  $k$  for all  $2 \leq k \leq \left\lfloor \frac{g+3}{2} \right\rfloor$ . However, for the finite set of reduced simplicial graphs of genus  $g$  this is not the case. For large enough  $g$  the upper bound is not sharp. This can be seen by combining Lemma 5.4 with the inequality that bounds the number of vertices in the beginning of the proof of Lemma 5.5.

We also define the Clifford index of a square-free module and a simplicial graph similarly as the Clifford index of a line bundle and a curve. That is, if  $M$  is a square-free module and  $d = \deg(M)$  and  $l(M) = r + 1$ , then we define  $\text{Cliff}(M) = d - 2r$ . For a 2-connected simplicial graph  $\Delta$  we define its Clifford index as

$$\text{Cliff}(\Delta) = \min\{\text{Cliff}(M) \mid M \text{ is a special effective module with } l(M) \geq 2 \text{ and } l(\omega_M) \geq 2\}.$$

As for curves we have the following proposition that implies that the Clifford index is  $\geq 0$ .

**Proposition 5.8** (Clifford's Theorem). *Let  $\Delta$  be a 2-connected simplicial graph of genus  $g \geq 2$ , and let  $M$  be a special effective  $k[\Delta]$ -module. If  $d = \deg(M)$  and  $l(M) = r + 1$ , then  $d \geq 2r$ .*

*Proof.* Suppose that  $M$  is a special effective module with  $l(M) = r + 1$ , and let  $s_1, \dots, s_{r+1} \in \tilde{H}_1(\Delta)$  be cycles corresponding to the global sections of  $M$ . We construct a new graph  $G$ , with a vertex  $v_i$  for each cycle  $s_i$  and an edge between each pair  $v_i, v_j$  if  $\dim \text{Supp}(s_i) \cap \text{Supp}(s_j) = 1$ . Since  $\Delta$  is 2-connected, we can use the same argument as in the proof of Lemma 5.2 to show that  $G$  is connected. Let  $e_1, \dots, e_n$  be an ordering of the edges of  $G$ , such that all the subgraphs  $H_i = e_1 \cup \dots \cup e_i$  of  $G$  are connected. Let  $M_i$  be the submodule of  $M$  generated by the cycles corresponding to the vertices in  $H_i$ . If  $e_1 = v_i v_j$ , then there are at least two vertices in

$\text{Supp}(s_i) \cap \text{Supp}(s_j)$  where the restriction from  $s_i$  and  $s_j$  are different, so the degree of  $M_1$  is at least 2. Suppose that  $\deg(M_i) = d_i$  and that there is a vertex in  $H_{i+1} \setminus H_i$  corresponding to a cycle  $s$ . Now we also have that there are at least two vertices in  $\text{Supp}(s)$  where the restriction of  $s$  is not in the span of the restriction of the cycles corresponding to the vertices in  $H_i$ . Hence  $\deg(M_{i+1}) \geq d_i + 2$ . It follows that the degree will increase by 2 for each vertex in the graph  $G$  except the first. In other words,  $d = \deg(M) = \deg(M_n) \geq 2v - 2 = 2(r + 1) - 2 = 2r$ .  $\square$

**Remark 5.9.** An interesting topic for further work was pointed out by F.O. Schreyer. When the graph is trivalent, the multiplication in the canonical module gives rise to a line configuration in  $\mathbf{P}^{g-1}$  in the following way. The projectification of the surjective multiplication maps  $\tilde{H}_{d-1}(\Delta) \cong (\omega_{k[\Delta]})_0 \xrightarrow{x_i} (\omega_{k[\Delta]})_{e_i} \cong \tilde{H}_{d-1}(\Delta; \Delta - \{i\})$  is a line in  $\mathbf{P}^{g-1}$ . The line configuration of the union of all these lines is a graph curve of arithmetic genus  $g$  which is canonically embedded. Furthermore, a  $g_d^1$  for the graph gives a projection of this curve to a  $\mathbf{P}^1$  of degree  $d$ . It is therefore natural to ask if the gonality of the graph and this curve is the same. Similar curves can also be made from more general 2-connected graphs, by choosing appropriate rational curves in the corresponding union of projective spaces. One interesting question is if there is a connection between the gonality or Clifford index of the graph and the minimal free resolution of this canonical curve as in Green’s conjecture for a canonical curve, which says that you can read the Clifford index of a curve of the graded betti numbers of the minimal free resolution. In many examples we made, this seems to be the case, but it seems very difficult to show this in general. This is first of all because the ideal of the canonical curve is not multigraded, and there is not a combinatorial way to describe the minimal free resolution. Secondly, the Clifford index of a graph is also not easily computed.

## REFERENCES

- [1] K. Baclawski. Canonical modules of partially ordered sets. *J. Algebra*, 83(1):1–5, 1983.
- [2] M. Baker and S. Norine. Riemann–Roch and Abel–Jacobi theory on a finite graph. *Advances in Math.*, 215(2):766–788, 2007.
- [3] D. Bayer and D. Eisenbud. Graph curves. *Advances in Math.*, 86:1–40, 1991.
- [4] W. Bruns and J. Herzog. *Cohen–Macaulay rings*. Cambridge University Press, 1998.
- [5] P.J. Cameron. *Combinatorics: Topics, Techniques, Algorithms*. Cambridge university press, 1994.
- [6] D. Eisenbud. *Commutative Algebra with a View Toward Algebraic Geometry*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, 1994.
- [7] E. Miller. The Alexander duality functors and local duality with monomial support. *J. Algebra*, 231(1):180–234, 2000.
- [8] E. Miller and B. Sturmfels. *Combinatorial Commutative Algebra*, volume 227 of *GTM*. Springer, 2000.
- [9] R.P. Stanley. *Combinatorics and commutative algebra*. Birkhauser, 1996.
- [10] K. Yanagawa. Alexander duality for Stanley–Reisner rings and square-free  $\mathbf{N}^n$ -graded modules. *J. Algebra*, 225(2):630–635, 2000.
- [11] K. Yanagawa. Derived category of squarefree modules and local cohomology with monomial ideal support. *J. Math. Soc. Japan*, 56(1):289–308, 2004.

MATEMATISK INSTITUTT, JOHS. BRUNSGT. 12, 5008 BERGEN  
*E-mail address:* `henning.lohne@math.uib.no`